Draft version February 2, 2008
Preprint typeset using LaTeX style emulateapj v. 6/22/04

arXiv:astro-ph/0503219v4 15 Jun 2005

BOUNDS ON EXPECTED BLACK HOLE SPINS IN INSPIRALING BINARIES

R. O'Shaughnessy[1], J. Kaplan[1], V. Kalogera[1], & K. Belczynski[2]

Draft version February 2, 2008

ABSTRACT

As a first step towards understanding the angular momentum evolution history of black holes in merging black-hole/neutron-star binaries, we perform population synthesis calculations to track the distribution of accretion histories of compact objects in such binaries. We find that there are three distinct processes which can possibly contribute to the black-hole spin magnitude: a birth spin for the black hole, imparted at either (i) the collapse of a massive progenitor star to a black hole or (ii) the accretion-induced collapse of a neutron star to a black hole; and (iii) an accretion spin-up when the already formed black hole [via (i) or (ii)] goes through an accretion episode (through an accretion disk or a common-envelope phase). Our results show that, with regard to accretion-induced spinup in merging BH-NS binaries [method (iii) above], only *accretion episodes associated with common-envelope phases and hypercritical accretion rates* occur in the formation history of merging black hole/neutron star binaries. Lacking unambiguous experimental information about BH birth spins [i.e., regarding the results of processes (i) and (ii)], we choose two fiducial values for the BH birth angular momentum parameter $a = J/M^2$, consistent with observations of (i) NS birth spins ($a \simeq 0$) and (ii) X-ray binaries ($a = 0.5$). Using these two fiducial values and a conservative upper bound on the specific angular momentum of accreted matter, we discuss the expected range of black hole spins in the binaries of interest. We conclude with comments on the significance of these results for ground-based gravitational-wave searches of inspiral signals from black hole binaries.

Subject headings: binaries:close — stars:evolution — stars:neutron – black hole physics – gravitational waves – relativity – X-rays:binaries

1. INTRODUCTION

Gravitational-wave astronomy is a rapidly-developing field, with many ground-based interferometric detectors (LIGO, VIRGO, GEO, TAMA) in place and taking data. These detectors are sensitive to (among other things) the waves emitted from the late stages of inspiral and merger of compact binaries with neutron stars (NS) and black holes (BH). Under certain conditions (as discussed in Apostolatos et al. 1994; Grandclement et al. 2003, 2004; Pan et al. 2004, and references therein) the BH spin in high-mass ratio binaries such as BH-NS binaries can significantly modulate the emitted inspiral waveform. While the more complex resulting waveforms can pose challenges for detection strategies, they also enable us to empirically determine the spin of rapidly-rotating black holes, if they exist. We undertake the present study of the expected BH spin distribution for two reasons: (i) to examine whether special search methods for spinning compact objects for optimal detection efficiency are necessary from an astrophysical point of view; (ii) to provide the first steps for a theoretical understanding of the origin and magnitude of BH spins in BH-NS binaries that will be useful when such BH spins are empirically constrained in the future.

In Sec. 2, we briefly describe the population synthesis calculations we use to explore a broad range of possible scenarios that lead to the formation of BH-NS binaries that can merge within a Hubble time (hereafter referred to as "merging"). We summarize and explain the common formation channels for these binaries and describe the mass accretion history associated with them. In Sec. 3, we review the observational evidence for BH birth spin drawn from observations of isolated NS and of X-ray binaries. Based on this evidence, we employ two fiducial choices for BH birth spin parameter $a = J/M^2$, i.e., $a = 0$ (nonspinning) or $a = 0.5$ (moderate spin). Given an understanding of the mass accretion history and a choice for the initial BH spin , we could in principle determine the final black hole spin if we knew the specific angular momentum of the material accreted by BH in these systems. This factor remains substantially uncertain. Thus, in Sec. 4 we use a *conservative* model for the specific angular momentum, combined with the distribution of accretion histories and birth spins, to limit the rate at which high-spin black holes occur in merging BH-NS binaries.

2. ACCRETION HISTORY OF BLACK HOLES IN BH-NS BINARIES

To generate and evolve stellar populations until double compact-object formation occurs, we use the *StarTrack* code first developed by Belczynski, Kalogera, and Bulik (2002) [hereafter BKB] and recently significantly updated and tested as described in detail in Belczynski et al. (2005). As with all binary population synthesis codes, *StarTrack* encapsulates the residual physical uncertainty in evolutionary processes of stars in a number of parameters, several of which can significantly affect the statistics of BH-NS mergers. In order to have a sample of merging BH-NS drawn from a large archive of astrophysically *a priori* plausible models, we generated an archive of accumulated results from a broad

---

[1] Northwestern University, Department of Physics and Astronomy, 2145 Sheridan Road, Evanston, IL 60208, USA

[2] Tombaugh Fellow, Department of Astronomy, New Mexico State University, Las Cruces, NM 88003

Electronic address: oshaughn, j-kaplan-1, vicky@northwestern.edu and kbelczyn@nmsu.edu

swath of plausible models for population synthesis. As described in O'Shaughnessy, Kalogera, and Belczynski (2004) [hereafter denoted OKB], the archive was generated using flat probability distributions for seven of the most significant parameters: three parameters describing the supernova kick magnitude distribution (assumed to consist of two maxwellian distributions with dispersions $\sigma_1 \in [0, 200]$km/s and $\sigma_2 \in [200, 1000]$km/s and relative weight of small to large kicks $s \in [0, 1]$); the effective common-envelope efficiency $\alpha\lambda \in [0, 1]$; the stellar wind strength $w \in [0, 1]$; the power-law index controlling the distribution of the mass of the companion relative to the mass of the primary $r \in [-3, 0]$; and the fraction of mass accreted (as opposed to lost from the binary) during non-conservative mass transfer episodes $f_a \in [0, 1]$. Unfortunately, as discussed in more detail in O'Shaughnessy et al. (2005b), we cannot succinctly describe the effect of these seven parameters on the formation channels and rates of BH-NS systems; physical predictions depend in a *highly correlated way on all seven parameters*. As in OKB, we fixed the remaining population synthesis model parameters to physically reasonable values, such as setting the maximum neutron star mass to $M_{NS,max} = 2M_\odot$ and the metallicity to solar metallicity $Z = 0.02$ (see parameters of model A in BKB for more details). From this archive we identified each merging BH-NS binary along with the details of its evolutionary history. Our results are summarized in Table 1.

TABLE 1
Statistics for evolutionary channels

Channel			n	AIC
(MT+)SNa	HCE(b→a)	(MT+)SNb	4310	3656
(MT+)SNb	HCE(a→b)	(MT+)SNa	543	543
(MT+)SNx		MT+SNy	319	0
(MT+)SNx		SNy	263	0
total:			5435	

Note. — This table summarizes all evolutionary channels followed by systems which formed merging BH-NS binaries in our simulations. The first three columns describe the evolutionary channel (where SNx indicates a supernovae of either *a* the primary or *b* the secondary, HCE indicates a hypercritical common envelope phase, and MT indicates a stable Roche-lobe-overflow mass transfer; parentheses indicate an optional feature of the channel); the fourth column provides the number of merging BH-NS binaries which passed through this channel; and the fifth lists the number which undergo accretion-induced-collapse (i.e., the BH forms during the common-envelope phase from an accreting neutron star). Most systems formed via a hypercritical common envelope (HCE) phase onto a compact object; this compact object is occasionally already BH, but usually is a NS which undergoes accretion-induced collapse (AIC) to a BH. A smaller number of systems (582 of 5435) do not undergo HCE at all and interact either weakly (through winds) or not at all between the two supernovae.

Compact binaries tight enough to merge through the emission of gravitational waves within the Hubble time have progenitors which usually interacted strongly (e.g. through mass transfer) earlier in their evolution. For example, a common mode of interaction is conventional mass transfer through Roche-lobe overflow and disk accretion. However, for merging BH-NS binaries, we find in our simulations that almost all of them have experienced a more dramatic phase of dynamically unstable mass transfer and common envelope (CE) evolution (see Table 1). During such phases it is expected that *hypercritical* accretion becomes possible. In this form of accretion (HCE), discussed in more detail in Brown (1995), Brown et al. (2000), and Fryer et al. (1996), a compact object spirals in through the envelope of its companion, rapidly accreting matter at highly super-Eddington (for photons), neutrino-cooled rates. As a result of this process, most of the companion's envelope is ejected, bringing the post-CE binary very close together. While the detailed accretion onto the compact object is complex and ill-understood, since we know the whole envelope will be lost during this phase, a straightforward application of the conservation of energy and mass during the quasicircular spiral-in process can determine the final orbit, the final binary masses, and the mass of the ejected material; see Appendix A of Belczynski et al. (2002) for details. Nevertheless we stress that our treatment of HCE is essentially a simple, semi-analytical model, which is by no means guaranteed to accurately describe the quantitative effect of the HCE phase.

More specifically, as described in Table 1, the vast majority (all but 582, out of 5435) of BH-NS binaries seen in our simulations form through an evolutionary channel that involves an HCE phase. In these channels, after an optional mass transfer phase, one star explodes, the compact remnant spirals through and strips the envelope of its companion, and then the companion explodes. Black holes form either immediately after the first core-collapse event or through the accretion-induced collapse of a NS that forms in the first supernova event. Regardless of the nature of the first compact object (BH or NS), it most typically experiences HCE evolution, and accretes a large amount of matter (potentially equal to its birth mass) during the brief HCE phase. Figure 1 shows the fraction of the initial compact object's mass $M_{\rm init}$ that is accreted during the HCE phase. Based on the sample we have, we estimate an upper bound on the fractional mass accreted as a function of the initial mass to be

$$(\Delta M/M)_{max} \approx 0.25 + 2M_\odot/M_{\rm init} \; . \tag{1}$$

This bound is valid for all channels.[3] For systems which form their BH immediately after the first supernova (i.e., no accretion-induced collapse, so $M_{\rm a,init} > 2M_\odot$) we find an empirical *lower* bound of

$$(\Delta M/M)_{max} \approx 0.05 + 0.6M_\odot/M_{\rm init} \; . \tag{2}$$

This latter bound may be understood as a requirement that the BH-NS system merge within the simulation time. Given that the second supernova will rarely further tighten the orbit, if the binary is to merge within 10 Gyr, then the orbit must be smaller than roughly $3.4R_\odot \left[(M_1/M_\odot)(M_2/M_\odot)(M_1+M_2)/M_\odot\right]^{1/4}$ after the common envelope phase. Using the common-envelope evolution equations presented in Appendix A of Belczynski et al. (2002) (with any choice for $\alpha_{\rm CE} \times \lambda$), we find this requirement approximately translates into the lower bound presented above.

Merging BH-NS can form through other channels, as shown in Table 1. However, these channels involve substantially less mass transfer onto the BH: for the HCE-related channels, the *smallest* mass increase

---

[3] We expect that systems which transfer more mass than this limit are brought so close that they merge during their common-envelope phase.



FIG. 1.— For systems that undergo HCE evolution (see Table 1), this plot shows $(M_{\rm final} - M_{\rm inital})/M_{\rm initial}$ versus $M_{\rm initial}$, where $M_{\rm initial}$ and $M_{\rm final}$ are the mass of the BH progenitor just after its supernovae and at the end of its evolution, respectively. In other words, they show the relative increase in mass of the compact object $\Delta M_{\rm HCE}/M$ due to hypercritical common envelope accretion. The upper solid line shows the empirical upper bound given in Eq. (1); the two vertical lines shows our assumed lower and upper limits on NS mass, $M = 1.3M_\odot$, $2M_\odot$.

was $0.22M_\odot$, to be contrasted with the *largest* disk-mediated mass transfer, $0.007M_\odot$ (i.e., from the channel SNx+MT+SNy). As described in Section 4, this incredibly small disk-mediated mass transfer cannot significantly modify the birth spin of the BH. Therefore, we neglect these channels in subsequent discussion.

Caveat: Branching ratios: Unfortunately, our branching ratios for various evolutionary channels are not significant. When exploring different population synthesis parameter combinations, we normalized so each population synthesis run contributes a roughly fixed number of merging NS-NS binaries to the archive. As a result, models associated with low NS-NS merger rates but high BH-NS merger rates should contribute disproportionately more merging BH-NS to our archive, and vice-versa. Moreover, even if each model contributed equally to the total number of BH-NS systems, our net numbers would then reflect the *mean branching ratios* over all population synthesis models considered, rather than the branching ratio of, e.g., the most common model. Finally, as discussed in O'Shaughnessy et al. (2005b), not all population synthesis models we considered are equally compatible with observations. Nonetheless given the current statistics, we fully expect that most BH-NS systems will form via a hypercritical common envelope phase.

3. BIRTH SPINS OF BH IN BH-NS BINARIES

No direct measurements of BH birth spins (or even BH spins) exist yet. A priori the BH birth spin cannot be constrained beyond the most fundamental level (i.e., to avoid naked singularities, we must have low angular momentum J: specifically, $a \equiv J/M^2 \leq 1$). For example, the collapse of a slightly-hypermassive NS rotating at breakup will, on dimensional grounds alone, produce a BH of spin of order half of maximal [$a \sim 0.5$] should it collapse; more realistic computations which allow for potential differential rotation support could push this value even higher (see, e.g., Duez et al. 2004; Shapiro 2002; Hawke et al. 2004).

Lacking unambiguous theoretical guidance for BH birth spins in merging BH-NS systems, and lacking direct observations of BH spin, we employ evidence for birth spin in similar systems to guide our choices for BH birth spin. On the one hand, on the basis of our understanding of NS birth spins and spinup in NS-NS binaries, we expect *low* BH birth spins ($a = 0$). However, on the basis of suggestive observations of X-ray binaries, we suspect *moderate* BH birth spins could occur ($a = 0.5$)

3.1. *BH birth spin estimates from NS observations*

The same processes that produce BH birth spin (i.e. core collapse and accretion-induced spinup) also determine the spins of the well-observed NS population. Observations of the NS population can thus potentially provide us with some constraints on BH birth spins.

BH birth spin from core collapse: We expect we can estimate the birth spins $a = J/M^2$ of BHs, and particularly low-mass BHs, through the estimated birth angular momenta J of young NS. For BHs and NSs of comparable mass, the collapse process should be nearly identical; therefore, the collapse product should have very similar J and M. Assuming no other process intervenes [for example, as discussed in Lindblom and Owen (2002), r-mode damping is not expected to significantly change the NS angular momentum of young NS], NS will spin down electromagnetically, and thus NS spins at birth can be estimated from the observed pulsar NS sample. In the past few years a number of studies of observed radio pulsars have estimated the NS spin periods at birth (see Lorimer 2004; Kramer 2003; Migliazzo et al. 2002, and references therein) in the range of $10 - 140\,$ms. These values are significantly slower than break-up spins (for typical NS equations of state) and correspond to $a \simeq 0.005 - 0.02$ (assuming a NS radius of 10 km and rigid rotation).

BH birth spin from accretion-induced collapse: Our simulations show that a significant fraction (i.e., 4199 out of 5435 binaries) of the BH in merging BH-NS systems were originally NS that experienced HCE and collapsed into BH. Therefore birth spins for this BH class are related to the spin-up of NS during a CE phase. Our current understanding of this spin-up process is quite limited. However, we can obtain guidance from the mildly recycled pulsars in known double neutron star binaries. These pulsars are believed to have been spun up during a CE phase (where collapse to a BH was avoided though) and it is evident that they are not spinning close to break-up: the fastest known pulsar in a double NS binary is PSR J0737-3039A spinning at $\simeq$ 20 ms or else having $a = 0.01$ (Burgay et al. 2004). Therefore post-CE NS spins appear to correspond to a no larger than $a = 0.01$.

Based on the observational considerations discussed above, we would expect the birth BH spin to be negligible (i.e., $a \lesssim 0.03$). However, a number of uncertainties in our modeling suggest that very different values of a could be equally plausible.

Uncertainties in collapse model: For black holes that form from the supernova of a massive star, we assume that the birth spin of these BHs are comparable to the birth spin of comparable-mass NSs, and we then assume that those NS birth spins may be estimated from electromagnetic (pulsar) spindown from present-day pulsars. However, most of these low-mass black holes form through fallback of post-supernova ejecta onto the hole; if the post-supernova fallback material carries significant angular momentum and can spin up the nascent BH to high values of a, even if the protoneutron star is not spinning with a period shorter than $\simeq 10\,\mathrm{ms}$. Further, if young protoneutron stars can spin down rapidly through other mechanisms, such as r-modes (Lindblom and Owen 2002), straightforward electromagnetic-based extrapolations may significantly underestimate the protoneutron star spin. If either of these mechanisms took place, then the birth BH spin could be significantly larger than the naive estimate we outlined above.

Uncertainties in HCE model: The known pulsars in double NS binaries have avoided accretion-induced collapse into BH. Those that do not avoid this collapse must accrete a higher amount of mass and therefore it is still possible that they get spun-up to shorter spin periods and therefore lead to BH formed with higher a values than expected.

Given these uncertainties, we cannot be sure whether the relevant BH birth spins are negligible (i.e., $a = 0$) or large, comparable to the break-up spin of a NS (i.e., $a \sim 0.5$).

3.2. *Black hole Spins in X-Ray Binaries*

Black hole X-ray binaries (XRBs) offer the possibility of *directly* measuring the spin of a compact object, since they involve a highly relativistic accretion flow in the strong field of a BH. Two techniques are prominent in the literature: interpreting quasiperiodic oscillations and fitting iron line profiles.

Quasiperiodic oscillations are believed to be modes of the inner, highly relativistic regions of the BH accretion disk. As such, their frequency is expected to be related to the Kepler frequency of the inner edge of the disk, which in turn is intimately connected with the properties of the black hole. Measurements of high-frequency QPOs that are inconsistent with the innermost stable circular orbit of a nonspinning black hole, as by Strohmayer (2001) and Remillard et al. (2002), have been interpreted as evidence for BH spin. Moreover, evidence suggests the spin suggested by these measurements is *not* merely a relic of past accretion. Unfortunately, QPO frequencies cannot be unambiguously translated to black hole angular momenta; their interpretation depends strongly on disk mode modelling (see, e.g. Torok et al. 2005; Rezolla et al. 2003), with results that vary from $a \approx 0.1$ to $a \approx 1$ depending on the assumptions used. However, even if the QPO interpretation is correct, the associated BH spin estimates need not reflect BH birth spins; such spins are most probably reached via long-term disk accretion, a process that is not relevant to BHs in BH-NS merging binaries.

Iron line profiles potentially offer in principle a more direct probe into the inner disk rotation profile and, thus, the BH spin. The iron line can show strong Doppler shifts (due to rotation) and redshifts (due to strong gravity); in particular, very rapidly spinning BHs should show asymmetric red "tails". While pioneered for use with AGN, this same technique has been applied to BH XRBs (see, e.g., Miller et al. 2002, 2004a,b). We treat the results, several claims of *extreme* spins $a > 0.8$, with great caution however. On the one hand, these systems are strongly model dependent, and many physically relevant details (e.g., proper accounting of light bending near the hole; see Beckwith and Done 2004) have yet to be included in the models. On the other hand, just like in the case of QPO interpretations, the estimated spins may be entirely due to the disk accretion in the XRB and therefore unrelated to the birth BH spin.

3.3. *Birth Black Hole Spins Adopted*

To summarize, based on observations of isolated NS we believe the birth spins of NS – and thus of low-mass BHs – to be small. On the other hand, observations of X-ray binaries (QPOs and iron lines) suggests that higher-mass BHs ($M > 5M_\odot$) could be born with moderate spin. To allow for this substantial uncertainty, in what follows we will consider the implications of two models for the birth BH spin: $a = 0$ and $a = 0.5$.

4. MASS ACCRETION AND SPINUP

Figure 1 demonstrates that the black hole accretes a significant fraction of its mass during the HCE phase. Since that matter probably carries some amount of angular momentum, the black hole could conceivably spin up significantly during the HCE phase. Unfortunately, the details of HCE accretion – particularly at the fine level of detail needed to resolve the amount of angular momentum advected onto the compact object – are not understood.

Lacking a quantitatively sound choice for the specific angular momentum, we argue instead that the specific angular momentum accreted should be *at most* the angular momentum of the marginally stable equatorial particle orbit. This model, equivalent to assuming that the accretion proceeds through a thin disk, implies that the final black hole spin depends only on the total amount of matter accreted and the initial black hole spin J (or, as commonly denoted, $a = J/M^2$); as derived initially by Bardeen (1970) (see also Thorne 1974), for a black hole which is initially nonspinning with mass M_i, the BH spin parameter a is given by the following expression $a = a_B(M, M_i)$:

$$a_B(M, M_i) = \left(\frac{2}{3}\right)^{1/2} \frac{M_i}{M} \left[4 - \left(\frac{18M_i^2}{M^2} - 2\right)^{1/2}\right]. \quad (3)$$

We consider this estimate to represent *an upper limit* to the BH a, since we do not expect that accretion in a HCE event will occur through a thin disk. Instead, we expect that on almost all scales the accretion flow will be nearly radial.

4.1. *Accretion onto nonspinning holes*

We use our results on compact object masses at the onset and the end of HCE to estimate a under the thin disk assumption. We consider two cases: (i) all mass accreted during the CE phases contributes to spinup and (ii) if accretion-induced collapse (AIC) occurs, only the fraction of the mass accreted after AIC contributes to spinup.

All mass contributes to spinup: As demonstrated in the top panel in Fig. 2, because of the substantial amount of mass accreted during HCE phases, nearly all BHs could conceivably spin up to very large a; a histogram of the values of a we observe would be highly concentrated near $a \sim 1$. In particular, many low-mass BHs (i.e, those which formed via AIC during an HCE phase) have accreted of order half their final mass, and can potentially



FIG. 2.— For systems that undergo HCE evolution (i.e., the "standard" or "reversed" channels presented in Table 1, this plot shows the BH spin expected assuming the HCE accretion occurs through a thin equatorial corotating disk: points are $(M_{a,\mathrm{final}}, a(\Delta M))$ where a is computed using the Bardeen formula [Eq. (3)] assuming an initially nonrotating hole. The two panels show the results assuming (i) all the mass that is accreted contributes to spinup and (ii) only the mass accreted after AIC of the NS into a BH. The solid curves (shown only for $M > 4$) show the empirical upper bound presented in Eq. (4) for nonspinning newborn BH.

spin up to of order $a \sim 1$. Further, *every* system that undergoes HCE evolution accretes enough mass to spin up significantly ($a > 0.2$). In this case, the BH-NS binaries with the largest spins (i.e., which have $a \simeq 1$) have mass ratios between $2:1$ to $3.5:1$.

However, for NS which undergo AIC, this approach approach will strongly overestimate the maximum attainable spin, since it assumes disk accretion at the schwarzchild radius, even before the BH forms.

Only post-AIC mass contributes to spinup: Since current astrophysical evidence suggests that NS do not get strongly spun-up during CE events, we also consider the possibility that only the mass accreted *after* the BH is formed through AIC contributes to its spin-up. (Those BH which form through direct collapse are treated as in the previous case.) In this case, summarized in the bottom panel in Fig. 2, we find a much broader distribution of possible a values (i.e., a histogram of possible a outcomes would be more nearly flat, from $a = 0$ to $a = 1$). In this case, the systems with the largest spins have mass ratio near $3.5:1$.

Both models have the same behavior for systems with large final BH masses ($M_{\rm bh,f} > 4M_\odot$). Good empirical lower and upper bounds for a for $M_{\rm bh,f} > 4M_\odot$ follows directly from the mass-accretion bounds Eqs. (1,2):

$$a < a_B\left(M_{bh,f}, \frac{4M_{bh,f}}{5} - 8M_\odot/5\right) \qquad (4)$$

$$a > a_B\left(M_{bh,f}, 0.95M_{bh,f} - 0.57M_\odot\right) \qquad (5)$$

where $M_{bh,f}$ and $M_{bh,i}$ are the BH final and birth masses, respectively.



FIG. 3.— As Fig. 2, but on a different vertical scale and assuming an initially *rotating* hole: $a = 0.5$. The solid curves (shown only for $M > 4$) show the empirical upper bound presented in Eq. (7).

4.2. *Accretion onto spinning holes*

Black holes which are initially spinning present a more complex accretion challenge in principle: the BH spin need not be aligned with the disk angular momentum axis, so the accreted material could just as well *spin down* as spin up the hole. However, following the same process we used to circumvent the the (substantial) uncertainties we addressed earlier, we limit attention to *bounding the BH spin.* The most conservative bound is obtained by assuming a corotating equatorial disk.

Bardeen's formula [Eq. (3)] applies equally well to BHs which are initially spinning, if M_i is chosen so the true initial black hole mass $M_{bh,i}$ and spin a_i satisfy $a_B(M_{bh,i}) = a_i$. To be concrete, if the birth BH spin is $a_i = 0.5$, then choosing

$$M_i = 0.84\, M_{bh,i} \qquad (6)$$

yields $a_B(M_{bh,i}, M_i) = 0.5$. When we repeat the analysis of the previous section, we find results summarized in Figure 3. Since Bardeen's spinup relation is monotonic, the empirical upper limit presented in Eq. (4) [for $a_i = 0$] translates directly into a corresponding spin limit when $a_i = 0.5$

$$a < a_B\left(M_{bh,f}, 0.84\times\left[\frac{4M_{bh,f}}{5} - 8M_\odot/5\right]\right) \quad . \qquad (7)$$

$$a > a_B\left(M_{bh,f}, 0.84\times\left[0.95M_{bh,f} - 0.57M_\odot\right]\right) \qquad (8)$$

Briefly, the results are qualitatively similar to the nonspinning case, but compressed in scale such that the minimum possible spin is $a = 0.5$ instead of $a = 0$. Since the construction of spin as a function of BH mass is similar, the spin upper bound also follows Eq. (4), with the Bardeen formula calculated choosing M_i so the initial black hole has spin $a = 0.5$.

4.3. *General Conclusions*

Our simulations have shown that almost all merging BH-NS binaries form through a hypercritical common envelope (HCE) phase, where they accrete a substantial fraction of their final mass. Lacking a quantitative model for the specific angular momentum accreted during HCE, we use the most pessimistic model: we assume the BH spins up through a thin equitorial disk, according to the Bardeen formula [Eq. (3)]. Even setting aside those systems with even greater physical uncertainties – those systems with final BH mass $M < 4M_{\odot}$, which form through accretion-induced collapse of the NS and whose final spin depends additionally on the details of the collapse and angular momentum transport onto the NS during that collapse – and even ignoring the possibility of BH birth spin, we find that large spins ($a \sim 1$) could easily be attained in principle from the HCE phase alone: with such a substantial fraction of its final mass transferred, the BH could easily spin up.

However, for BHs with $M > 4M_{\odot}$, significant spinup after birth becomes increasingly difficult, as larger black holes require ever larger (and more unlikely) mass transfers in order to spin up to large a. For example, assuming nonrotating initial BHs, the only BHs we found with spin $a > 0.8$ are those with final masses between $3M_{\odot}$ and $10M_{\odot}$; similarly, for BHs with $a = 0.5$ at birth, the only BHs we found with spin $a > 0.95$ are those with final masses in this same range.

5. BH SPIN AND DETECTION OF BH-NS INSPIRAL

In BH-NS binaries, the spin of the BH can, if large, have a significant effect on the late stages of inspiral and in particular the emitted gravitational waveforms (see Apostolatos et al. 1994; Apostolatos 1996; Grandclement et al. 2004; Pan et al. 2004; Buonanno et al. 2004, and references therein). As a result, in BH-NS binaries measurements of gravitational waves offer the potential to extract information about the spin. Furthermore, because of precession, such binaries will be much more reliably detectable, suffering less frequently from the effects of poor source orientation.

But these advantages come at a price. The more complex waveforms of spinning binaries are vastly more complicated to model. At present only ad-hoc models are considered computationally feasible (Pan et al. 2004), and even these methods involve substantial computational challenges, particularly at more extreme mass ratios.

Our calculations provide extremely conservative upper bounds on the BH spin in BH-NS binaries. These bounds tell us that employing spin need not pose such a dramatic dilemma, on astrophysical grounds: if we employ spin, we could require fewer computational resources than we might expect; and if we neglect it, we will not lose a dramatic fraction of potentially observable events.

5.1. *Mass ratio spin constraints and detection*

When constructing template banks for BH-NS inspiral a priori, parameter ranges are typically chosen to cover a very conservative parameter range; for example, in one of the few papers that presented results for variable mass ratio, Pan et al. (2004) built a template bank based on all binaries with masses in the range $M_{bh} \in [7, 12]M_{\odot}$ and $M_{ns} \in [1, 3]M_{\odot}$ and BH spins in the range $a \in [0, 1]$.

However, most of the templates in the template bank come from extreme-mass-ratio systems, involving a high-mass BH which we have just demonstrated is difficult to spin up. Therefore, a different template bank, which neglected astrophysically irrelevant systems, could perform the same search much more rapidly. Detailed computations of the computational savings we could obtain by moderating our template bank are far beyond the scope of this paper, and will depend sensitively on two poorly constrained parameters, the BH birth spin and the maximum NS mass (here, assumed $M = 2M_{\odot}$.) However, in an optimistic case we expect that a more judicious choice of templates could reduce the template bank size by a significant factor, cf. Sec. VI C of Pan et al.

5.2. *Limits on the decrease in event rate due to omitting BH spin*

For the purposes of detection, spinning templates are required only when nonspinning templates fail to have sufficient overlap with the physical signal to guarantee most inspiral events are detected. Loosely speaking, a nonspinning template fails to mimic the effect of *precession* on the waveform. Because of spin-orbit coupling, angular momentum will be exchanged, causing the orbital plane to precess around the total angular momentum. Therefore, spinning templates are needed to detect those systems where the effects of precession are strongest – namely, those with (i) strong BH spins $a \approx 1$ that are (ii) strongly misaligned with the orbital angular momentum (see, e.g., Fig. 2 in Grandclement et al. 2004).

Grandclement et al. (2004) have quantitatively examined the expected overlap between nonspinning templates and the expected signal from binaries in which the BH spin magnitude and orientation are independently varied (see their Figure 2), assuming BH-NS binaries with $M_{bh} = 10M_{\odot}$ and $M_{ns} = 1.4M_{\odot}$. They then further convolved this function with an expected spin orientation distribution to derive an estimate for the fraction of events that could be seen that will be seen, as a function of spin magnitude of the BH (see their Figure 3).[4] They find that even if all BH were spinning maximally, only about 30% of potentially-detectable BH-NS mergers will be missed.

The Grandclement et al. results, however, are not directly applicable in our circumstances. Spin-dependent modulations depend significantly on mass ratio, a parameter that is varied in our BH-NS binaries but fixed for those of Grandclement et al. Nonetheless, if we employ their relation between (i) the probability of detecting a detectable source when ignoring spin and (ii) the BH spin magnitude of that source, then we find when we convolve with our Monte Carlo sample of BH spins – which, recall, was designed to produce the largest plausible spin values, given known mass transfer – that we would lose at most 30% of events in extreme cases (i.e., high BH birth spin or all HCE mass accreted onto NS) and slightly less (25%) in more conservative cases (i.e., mass before AIC does not contribute to spin). Briefly, searches which use nonspinning template banks will likely find most detectable inspiral signals. Practically speaking, given the considerable uncertainty in the underlying event rate itself, upper limits produced using precise templates will have only marginally more astrophysical impact than upper limits produced using nonspinning template banks. Of course, for *parameter estimation*, correct templates are essential in order to extract the BH spin, and thus to constrain its accretion history.

---

[4] Specifically, they plot the average of the cube of the fitting factor versus the BH spin a.

6. SUMMARY AND FUTURE DIRECTIONS

In this paper we perform population synthesis calculations to track the history of systems which evolve into merging BH-NS binaries, demonstrating that these objects form through only a few channels, almost exclusively involving a hypercritical common envelope (HCE) phase and often involving accretion induced collapse of a NS into a BH. We show that a significant amount of mass is typically accreted during this HCE phase. Using *an upper limit based on thin-disk accretion*, we find the BHs accrete enough matter to potentially spin up to large a values, independent of the BH birth spin. Finally, we note the observed sample of neutron stars suggests the birth spins of the black holes in these binaries should be small; therefore, the BH spins in BH-NS binaries should arise almost entirely from the matter they accrete in a HCE phase. Unfortunately, the HCE phase is poorly understood. While our result is a first step of a program to better understand the expected BH spins for those systems LIGO and other gravitational-wave detectors could observe, more simulations of these processes are needed to obtain firmer conclusions.

Further, based on our rough understanding of the range of expected spin magnitude, and using the expected effect of spin on GW detection drawn from Grandclement et al. (2004), we demonstrated that neglecting spin in GW searches should only moderately reduce the detection rate. However, our analysis was based on the work of Grandclement et al. (2004), which like many papers in the gravitational-wave literature assumed all BHs in BH-NS binaries had $M = 10M_{\odot}$. In fact, based on our calculations we expect (i) the mass ratio can vary, and should be biased towards larger values (i.e., closer to 3 : 1 than 10 : 1.4); and (ii) the sample of BH-NS binaries should show strong correlations between mass ratio and spin magnitude, because higher-mass BHs are harder to spin up. We hope to undertake a more thorough Monte Carlo study of the effect of various expected distributions for BH spin magnitude, spin tilt, and BH-NS mass ratio in a future paper.

Finally we note that in our simulations we consider only BH-NS in the Galactic field; we do not account for any stellar interactions relevant to centers of globular clusters and similar dense environments. If BH-NS binaries form in significant numbers in cluster centers, then their spin properties could be entirely unrelated to the analysis presented in this study.

We thank Ron Taam, Fred Rasio, and Stu Shapiro for helpful discussions. We particularly thank our anonymous referee for extensive comments and suggestions regarding the XRB literature. This work is partially supported by NSF Gravitational Physics grants PHYS-0121416 and PHYS-0353111, a David and Lucile Packard Foundation Fellowship in Science and Engineering, and a Cottrell Scholar Award from the Research Corporation to VK. VK also thanks the Aspen Center for Physics for its hospitality while working on this paper.

REFERENCES

Apostolatos, T. A., Cutler, C. , Sussman, G. J., and Thorne, K. S. 1994, Phys. Rev. D, 49, 6274
Apostolatos, T. A. 1996, Phys. Rev. D, 54, 2438
Bardeen, J. M. 1970, Nature, 226, 64.
Beckwith, K., and Done, C. 2004, MNRAS, 352, 353.
Belczynski, K., Kalogera, V., and Bulik, T. 2002, ApJ, 572, 407
Belczynski, K., Kalogera, V., Taam, R., Rasio, F., Zezas, A., Bulik, T., Maccarone, T., Ivanova, N. 2005, ApJ, to be submitted
Brown, G. E., Lee, C.-H., and Bethe, H. A. 2000, ApJ, 541, 918.
Brown, G. E. 1995, ApJ, 440, 270.
Buonanno, A., Chen, Y., Pan, Y, and Vallisneri, M. 2004, Phys. Rev. D, 70, 104003
Burgay, M. et al. 2003, Nature, 426, 531
Duez, M, Liu, Y. T., Shapiro, S, and Stephens, gr-qc/0402502
Kramer, M., et al. 2003, MNRAS, 342, 1299.
Fryer, C. L., Benz, W., and Herant, M. 1996, ApJ, 460, 801.
Grandclement, P., Kalogera, V., Vecchio, A. 2003, Phys. Rev. D, 67, 042003
Grandclement, P., Ihm, I, Kalogera, V., and Belczynski, K. 2004, Phys. Rev. D, 69, 102002
Hawke, I., et al. 2004, Phys. Rev. D, 71, 024035 gr-qc/0403029
Lindblom, L., and Owen, B. 2002, Phys. Rev. D, 65, 063006
Lorimer, D. 2004, Aspen Winter Conference on Astrophysics (proceedings in press; available from `www.astro.northwestern.edu/AspenW04/program.html`)
Remillard, R., et al. 2002, ApJ, 580, 1030
Migliazzo et al. 2002, in *Neutron Stars in Supernova Remnants*, ASP Conference Series Vol. 271, 57
Miller et al. 2002, ApJ, 570, L69
Miller et al. 2004, ApJ, 601, 45
Miller et al. 2004, ApJ, 601, L131
O'Shaughnessy, R., Kalogera, V. , and Belczynsi, K. 2005, ApJ, 620, 385
O'Shaughnessy, R., Kim, C., Fragos, T., Kalogera, V., and Belczynski, K. 2005, ApJ, submitted
Pan, Y, Buonanno, A., Chen, Y., and Vallisneri, M. 2004, Phys. Rev. D, 69, 104017
Portegies-Zwart, S. F. & Yungelson, L. R. 1998, A& A, 332, 173
Rezolla, L., et al. 2003, MNRAS, 344, L34.
Shapiro, S. and Shibata, M. 2002, ApJ, 577, 904.
Strohmayer, T. E. 2001, ApJ, 552, L49
Thorne, K. S. 1974, ApJ, 191, 507
Torok, G., Abramowicz, M. A., Kluzniak, W., and Stuchlik, Z. 2005, A&A 436, 1